

**Table 2.1.10: Catch-up schedule for 13vPCV\* (Prevenar 13) for Indigenous children residing in the Northern Territory, Queensland, South Australia or Western Australia ONLY, who do not have any medical condition(s) associated with an increased risk of invasive pneumococcal disease (IPD), aged <5 years**

| Number of doses given previously | Age at presentation | Age when previous dose of any PCV <sup>†</sup> was given |             |            | Recommendations   |  |
|----------------------------------|---------------------|--|-------------|------------|---|--|
|                                  |                     | 1st dose   | 2nd dose    | 3rd dose   | Number of further primary dose(s) required <sup>‡</sup> | Number of booster 13vPCV doses required <sup>§</sup> |
| No previous doses                | <7 months           | –  | –           | –          | 3   | 1  |
|                                  | 7–11 months         | –  | –           | –          | 2   | 1  |
|                                  | 12–23 months        | –  | –           | –          | 1   | 1  |
|                                  | 24–59 months        | –  | –           | –          | 1   | Not needed   |
| 1 previous dose                  | <7 months           | <7 months  | –           | –          | 2   | 1  |
|                                  |                     | 7–11 months  | –           | –          | 2   | 1  |
|                                  | 7–11 months         | <7 months  | –           | –          | 1   | 1  |
|                                  |                     | 7–11 months  | –           | –          | 1   | 1  |
|                                  | 12–23 months        | <12 months   | –           | –          | 1   | 1  |
|                                  |                     | ≥12 months   | –           | –          | Not needed  | 1  |
| 24–59 months                     | <12 months          | –  | –           | 1          | Not needed  |  |
|                                  | ≥12 months          | –  | –           | Not needed | Not needed  |  |
| 2 previous doses                 | <12 months          | <7 months  | <12 months  | –          | 1   | 1  |
|                                  |                     | 7–11 months  | 7–11 months | –          | Not needed  | 1  |
|                                  | 12–23 months        | <7 months  | <12 months  | –          | 1   | 1  |
|                                  |                     | 7–11 months  | Any age     | –          | Not needed  | 1  |
|                                  |                     | ≥12 months   | ≥12 months  | –          | Not needed  | Not needed   |
|                                  | 24–59 months        | Any age  | Any age     | –          | Not needed  | Not needed   |
| 3 previous doses                 | 7–11 months         | Any age  | Any age     | Any age    | Not needed  | 1  |
|                                  | 12–23 months        | <7 months  | <12 months  | Any age    | Not needed  | 1  |
|                                  |                     |  | ≥12 months  | ≥12 months | Not needed  | Not needed   |
|                                  | 7–11 months         | 7–11 months  | ≥12 months  | Not needed | Not needed  |  |
| 24–59 months                     | Any age             | Any age  | Any age     | Not needed | Not needed  |  |

\* If 13vPCV is not available, and 10vPCV is being used for all/any children, 10vPCV is recommended in a 4-dose schedule for infants (i.e. at ages 2, 4, 6 and 12–18 months). If catch-up is required for 10vPCV, vaccination can be done according to the information provided in this Table. (See also 4.13 *Pneumococcal disease*.)

† Prior PCV doses may have been given as 7vPCV (e.g. from overseas), 10vPCV or 13vPCV. 13vPCV should be used as the vaccine formulation for further catch-up doses required, regardless of which formulation of PCV the child received previously.

‡ Recommended interval between primary doses for catch-up is 1–2 months. Where possible, it is recommended to align doses with the standard schedule points at 4 months and 6 months of age for infants aged <7 months. The minimum interval between dose(s) is 1 month if aged <12 months, and 2 months if aged ≥12 months.

§ A minimum interval of 2 months is required after the last dose of 13vPCV in the primary course.