

Table 2.1.11: Catch-up schedule for 13vPCV (Prevenar 13) and 23vPPV (Pneumovax 23) in children with a medical condition(s) associated with an increased risk of invasive pneumococcal disease (IPD),* presenting at age <2 years

For children with a medical condition(s) associated with an increased risk of IPD presenting at age ≥2 years, see recommendations in 4.13 Pneumococcal disease and Table 4.13.2.†

Number of doses given previously	Age at presentation	Age when previous dose of any PCV‡ was given			Recommendations		
		1st dose	2nd dose	3rd dose	Number of further primary dose(s) of 13vPCV required§	Number of booster doses of 13vPCV required at age ≥12 months¶	Number of doses of 23vPPV required at age 4–5 years
No previous doses	<7 months	–	–	–	3	1	1
	7–11 months	–	–	–	2	1	1
	12–23 months	–	–	–	1	1	1
1 previous dose	<7 months	<7 months	–	–	2	1	1
	7–11 months	<7 months	–	–	2	1	1
		7–11 months	–	–	1	1	1
	12–23 months	<12 months	–	–	1	1	1
		≥12 months	–	–	Not needed	1	1
2 previous doses	<12 months	<7 months	<12 months	–	1	1	1
		7–11 months	7–11 months	–	Not needed	1	1
	12–23 months	<7 months	<12 months	–	1	1	1
			≥12 months	–	Not needed	1	1
		7–11 months	Any age	–	Not needed	1	1
		≥12 months	≥12 months	–	Not needed	Not needed	1
3 previous doses	7–11 months	Any age	Any age	Any age	Not needed	1	1
	12–23 months	<7 months	<12 months	Any age	Not needed	1	1
			≥12 months	≥12 months	Not needed	Not needed	1
		7–11 months	7–11 months	≥12 months	Not needed	Not needed	1

* Refer to List 4.13.1 in 4.13 *Pneumococcal disease* for the list of specified conditions.

† Recommendations for vaccination of haematopoietic stem cell transplant (HSCT) recipients differ; see Table 3.3.3 *Recommendations for revaccination following HSCT in children and adults, irrespective of previous immunisation history*.

‡ Prior PCV doses may have been given as 7vPCV (e.g. from overseas), 10vPCV or 13vPCV. 13vPCV should be used as the vaccine formulation for further catch-up doses required, regardless of which formulation of PCV the child received previously.

§ Recommended interval between primary doses for catch-up is 1–2 months. Where possible, it is recommended to align doses with the standard schedule points at 4 months and 6 months of age for infants aged <7 months. The minimum interval between dose(s) is 1 month if age <12 months, and 2 months if age ≥12 months.

¶ The booster dose of 13vPCV should be given at the earliest opportunity after the child reaches 12 months of age, but a minimum interval of 2 months is required after the last dose of 13vPCV in the primary course.