

Table 2.1.9: Catch-up schedule for 13vPCV (Prevenar 13) for non-Indigenous children, and Indigenous children residing in the Australian Capital Territory, New South Wales, Tasmania and Victoria, who do not have any medical condition(s) associated with an increased risk of invasive pneumococcal disease (IPD), aged <5 years

| Number of doses given previously | Age at presentation | Age when previous dose of any PCV* was given | | | Number of further dose(s) required [†] |
|----------------------------------|---------------------|--|-------------|----------|---|
| | | 1st dose | 2nd dose | 3rd dose | |
| No previous doses | <7 months | – | – | – | 3 |
| | 7–11 months | – | – | – | 2 |
| | 12–59 months | – | – | – | 1 |
| 1 previous dose | <7 months | <7 months | – | – | 2 |
| | 7–11 months | <7 months | – | – | 2 |
| | | 7–11 months | – | – | 1 |
| | 12–59 months | <12 months | – | – | 1 |
| | | ≥12 months | – | – | Not needed |
| 2 previous doses | <12 months | <7 months | <12 months | – | 1 |
| | | 7–11 months | 7–11 months | – | Not needed |
| | 12–23 months | <7 months | <12 months | – | 1 |
| | | | ≥12 months | – | Not needed |
| | | | ≥7 months | Any age | – |
| | 24–59 months | Any age | Any age | – | Not needed |
| 3 previous doses | 7–59 months | Any age | Any age | Any age | Not needed |

* Prior PCV doses may have been given as 7vPCV (e.g. from overseas), 10vPCV or 13vPCV. Use 13vPCV as the vaccine formulation for further catch-up doses required, regardless of which formulation of PCV the child received previously.

† Recommended interval between primary doses for catch-up is 1–2 months. Where possible, it is recommended to align doses with the standard schedule points at 4 months and 6 months of age for infants aged <7 months. The minimum interval between dose(s) is 1 month if aged <12 months, and 2 months if aged ≥12 months.