

**Table 4.13.2: Recommendations for pneumococcal vaccination for children aged 2–5 years with a medical condition(s) associated with an increased risk of invasive pneumococcal disease (IPD) (see List 4.13.1, Categories A and B)\***

Vaccination history <sup>†</sup>		Recommendations	
Primary course of any pneumococcal conjugate vaccine	Supplementary/booster dose of 13vPCV (at age ≥12 months)	Number of further 13vPCV dose(s) required for children aged 2–5 years	Single 23vPPV dose required at age 4–5 years <sup>‡</sup>
Completed	Received	0	Yes
Completed	Not received	1	Yes
Not completed	Not received	2 <sup>§</sup>	Yes

\* HSCT recipients require 3 doses of 13vPCV post transplantation, followed by 23vPPV, irrespective of previous vaccine doses received (see Table 3.3.3 *Recommendations for revaccination following HSCT in children and adults, irrespective of previous immunisation history*).

† See also 2.1.5 *Catch-up*, Table 2.1.11 *Catch-up schedule for 13vPCV (Prevenar 13) and 23vPPV (Pneumovax 23) in children with a medical condition(s) associated with an increased risk of IPD, presenting at age <2 years*. Note: this table primarily refers to 13vPCV, but could apply to use of 10vPCV if the former was not available.

‡ At least 2 months after the last dose of PCV, whichever is later.

§ Minimum interval between the 2 doses should be 2 months.